



OPEN ACCESS

Vaping and harm in young people: umbrella review

Su Golder,¹ Greg Hartwell ,² Lily M Barnett,³ Sophie G Nash,³ Mark Petticrew,² Rebecca E Glover

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/tc-2024-059219>).

¹Department of Health Sciences, University of York, York, UK

²Public Health, Environments and Society, Faculty of Public Health and Policy, London School of Hygiene & Tropical Medicine, London, UK

³Royal Free London NHS Foundation Trust, London, UK

Correspondence to

Dr Greg Hartwell;
gregory.hartwell@lshtm.ac.uk

SG and GH are joint first authors.

Received 12 December 2024
Accepted 10 June 2025

ABSTRACT

Objective To appraise and synthesise the evidence on short and longer term harms of vaping in young people.

Data sources KSR Evidence (systematic reviews); Medline, Embase, and PsycInfo (umbrella reviews); and reference screening.

Study selection Systematic and umbrella reviews evaluating any potential harms from e-cigarettes in young people.

Data synthesis Searches identified 56 reviews for inclusion from 384 unique articles. A consistent significant association between vaping and smoking initiation was found, supporting a causal relationship, with pooled ORs of 1.50–26.01 (21 systematic reviews), most of which suggested that young people using e-cigarettes are about three times more likely than those not using them to initiate smoking. Five systematic reviews demonstrated a substantial association between e-cigarettes and substance use, with pooled ORs of 2.72–6.04 for marijuana, 4.50–6.67 for alcohol and 4.51–6.73 for binge drinking. Asthma was the most common respiratory outcome, with consistent associations (ORs: 1.20–1.36 for diagnosis and 1.44 for exacerbation). Three systematic reviews found associations between vaping and suicidal outcomes, and six included investigation of injuries, predominantly documenting explosion incidents. Significant associations between vaping and other harmful outcomes included pneumonia, bronchitis, lower total sperm counts, dizziness, headaches, migraines and oral health harms, but this evidence was largely derived from limited surveys or case series/reports.

Conclusions This study found that there were consistent associations between vaping and subsequent smoking, marijuana use, alcohol use, asthma, cough, injuries and mental health outcomes. The findings support the implementation of policy measures to restrict sales and marketing of e-cigarettes to young people.

Trial registration number PROSPERO CRD42024569728.

INTRODUCTION

Youth vaping has increased markedly in recent years. The US Centers for Disease Control and Prevention estimated in 2024 that 5.9% of American middle and high schoolers were currently using e-cigarettes,¹ and the WHO describes the rise in children vaping around the world as "alarming".² Across WHO Europe region countries, e-cigarette use among those aged 15–16 years has been shown to range from 5.5% to 41%.³ For adults, e-cigarettes have been adopted and advocated as a smoking cessation tool by public health bodies in the UK and some other countries.^{2,4} However, the WHO is concerned about vapes being "aggressively

marketed" to young people and has called on governments around the world to strengthen their vaping policies.^{2,5}

The harm or safety of vaping products is thus a timely question for tobacco control internationally, and evidence synthesis is one lever to improve the evidence base underpinning policy in this area. There is ongoing concern about the harms of vaping in young people, especially whether e-cigarettes increase the risk of smoking initiation, a key area where a contemporary synthesis of evidence would be valuable. Although four umbrella reviews exist, these do not focus on young people alone or do so only for a narrow set of outcomes (such as smoking initiation).^{6–9} No umbrella review to date has specifically focused on young people and considered a full range of potential physical and mental health impacts in this age group. We therefore conducted an umbrella review to identify systematic reviews on the acute and longer-term harms of vaping in young people.

METHODS

This overview of reviews is reported in line with PRIOR (preferred reporting items for overviews of reviews: online supplemental file 2)¹⁰ and followed a prespecified protocol (PROSPERO CRD42024569728). The inclusion criteria were broad to provide an overall understanding of the health implications of vaping among young people (table 1).

Search methods

KSR Evidence database was searched on 4 July 2024, with update searches on 21 November 2024. KSR Evidence aims to include all systematic reviews, meta-analyses and health technology assessment reports published in the past decade and includes sensitive searches of several databases (including Medline, Embase, CINAHL, PsycInfo and the Cochrane Database of Systematic Reviews).¹¹ We also checked the references of included systematic reviews for related systematic reviews.

We included umbrella reviews to identify further systematic reviews; findings from these umbrella reviews are reported separately to avoid duplication. As KSR Evidence does not include umbrella reviews, we conducted a separate search on 21 November 2024 on Medline, Embase and PsycInfo for these, limited to a publication date of 2015 onwards, reflecting the major shift in types of e-cigarette devices being used by young people at this time (specifically the dominance of rechargeable tank/reservoir models after this point, followed by the surge in disposable vapes),¹² to avoid including reviews focused solely on 'cigalikes' or other



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY. Published by BMJ Group.

To cite: Golder S, Hartwell G, Barnett LM, et al. *Tob Control* Epub ahead of print: [please include Day Month Year]. doi:10.1136/tc-2024-059219

Table 1 Inclusion and exclusion criteria for reviews on the harms of vaping in young people

	Inclusion criteria	Exclusion criteria
Population	Children, young adults, adolescents and youths (<25 years), including reviews with wider age ranges if separate analysis of young people was provided, and reviews of subgroups, such as disease or condition specific groups (eg, pregnant women) if <25 years	Adults aged >24 years. Animals or in vitro studies
Exposure	Use of e-cigarettes/vaping (legal or illegal), electronic nicotine delivery systems and electronic non-nicotine delivery systems	Marijuana vaping, aromatherapy vaping, traditional cigarettes and cigars, secondhand vaping and nicotine patches
Comparator(s)	None or any with no specific criteria. Comparators can be defined by the criteria of the reviews themselves (often young people who have never used e-cigarettes, or those who are not currently using them)	No exclusions
Outcome	Physical and mental health harms (respiratory outcomes including e-cigarette or vaping use associated lung injury, uptake of conventional cigarette smoking or uptake of substance use, cardiovascular outcomes, gastrointestinal outcomes, mental health outcomes including depression or anxiety, and any other negative outcome)	Cigarette quitting, positive benefits (such as reduced stress), pregnancy harms to fetus, accidental overdose or poisoning
Setting	All countries/settings	None
Study design	Systematic review, umbrella review	Scoping review, non-systematic literature review, commentary, editorial or letter, primary study (randomised controlled trial or observational study)
Restrictions	After 2014. English language (or translation available). All types of publication.	Before 2015. Non-English language paper where no translation was available

obsolete devices. Search results were deduplicated in EndNote (Clarivate).¹³ Title and abstract screening were undertaken independently by two reviewers in Covidence (Covidence AS)¹⁴ with disagreements resolved by discussion or with a third reviewer. Full text screening was also undertaken in Covidence by two reviewers, with disagreements resolved in the same way.

Data extraction

Data were extracted on study characteristics, methods and results of the included systematic reviews and umbrella reviews in Covidence (see online supplemental files). In some reviews, there was a wider age range than our inclusion criteria; in such instances, only the results relating to young people were extracted. One reviewer carried out data extraction which was checked by a second reviewer.

Methodological quality

The quality of the included systematic reviews was assessed by one reviewer using AMSTAR 2¹⁵ and then checked by a second reviewer. Study overlap was assessed by calculating corrected covered areas (CCA; 0–5%=slight; 6–10%=moderate; 11–15%=high; >15%=very high overlap; CCA=N-r/(rc)-r, where N is the number of times publications appeared in reviews, r is the number of rows and c is the number of columns; see online supplemental file).¹⁶ Forest plots were created using odds ratios (ORs) from meta-analyses reported in those reviews that included a meta-analysis.

Patient and public involvement

The Youth Advisory Panel of the UK Association for Young People's Health (AYPH) provided input on emerging results, implications and dissemination via a presentation to an online meeting of the panel by one research team member, and a discussion co-facilitated by an AYPH Youth Engagement and Partnership Project Manager.

RESULTS

The database searches identified 425 records (381 after duplicates removed) and an additional three records were found from

reference list checking (figure 1). After screening 384 unique records, 195 full text papers were examined, and 56 reviews were included. The main reason for exclusion at full text review stage was a lack of analysis of data relating to young people (see online supplemental file). The included systematic reviews were published between 2016 and 2024, with 84% (47/56) published from 2020 onwards. Results are presented according to health outcome (table 2).

E-cigarette use and smoking initiation and relapse

Twenty-one systematic reviews^{9,17–37} and four umbrella reviews^{6–9} (one study⁹ described itself as an umbrella review and systematic


**Figure 1** Flow diagram for included systematic reviews and umbrella reviews. SR, systematic review; YP, young people.

Table 2 Study characteristics

	No	%
Review types		
Umbrella reviews	4	7.1
Systematic reviews	52	92.9
Harm outcomes*		
Smoking initiation	24	42.9
Substance use	6	10.7
Asthma	8	14.3
Respiratory (excluding asthma)	10	17.9
Injuries	7	12.5
Mental health	9	16.1
Cardiovascular	3	5.4
Oral health	5	8.9
Fertility	2	3.6
Other (COVID, headache/migraine, vascular damage, renal)	4	7.1
Quality assessment		
High	1	1.8
Moderate	2	3.6
Low	14	25.0
Critically low	39	69.6

*Percentages sum to >100 because some reviews reported on multiple harms.

review) assessed the evidence that e-cigarette use increases the risk of conventional cigarette use (online supplemental file 1). Fourteen countries were represented in the included primary studies: USA, Canada, Mexico, UK, Germany, Netherlands, Taiwan, Finland, Romania, Switzerland, Belgium, South Korea, Thailand and Lebanon. Where the total number of participants included was reported, this ranged from 2084 to 779 937.

Smoking Initiation

Four umbrella reviews included systematic reviews assessing the association of e-cigarettes and conventional cigarettes.^{6–9} None was limited to young people. While our umbrella review identified 21 systematic reviews for inclusion, the four previous umbrella reviews included eight,⁷ three,⁹ two⁸ and one review,⁶

respectively. The included systematic reviews^{9 17 18 20 26 27 29 34 38} within these umbrella reviews were all captured and included in our review, except Zhong *et al*,³⁸ which only measured intentions to smoke, not actual smoking behaviours. All umbrella reviews found evidence that young people who use e-cigarettes are more likely than young people who do not to initiate cigarette smoking.

All 21 systematic reviews demonstrated an association between e-cigarette use and subsequent initiation of smoking independent of whether they assessed ever or current e-cigarette use with ever or current smoking. The reviews included 323 primary studies (90 unique), mostly longitudinal cohorts with follow-up durations of 4 months to 2 years; 46% (41/90) of the primary studies were included in multiple reviews ranging from 2 to 16 reviews (online supplemental table). The degree of overlap was high (CCA=13%).³⁹

ORs ranged from 1.50–26.01, with most reporting that young people using e-cigarettes are about three times more likely than those not using them to initiate smoking. Within individual reviews, there was heterogeneity in the magnitude of ORs of primary studies, but the consistent, statistically significant associations between e-cigarettes and smoking initiation are clear from the adjusted ORs presented in the 12 systematic reviews with pooled meta-analysis (online supplemental file 1 and figure 2). ORs were significant whether using adjusted or unadjusted values (factors adjusted for were mostly sex, age and socioeconomic status, with some results adjusted for other variables, such as psychosocial/behavioural risk factors and family/peer smoking). While all review authors concluded that the use of e-cigarettes was associated with an increased risk in smoking initiation, some questioned whether this was a causal effect.

Flavoured e-liquid and smoking initiation

Previous research has found that flavoured e-cigarettes are more popular among younger people,⁴⁰ but only one systematic review studied associations between the use of e-liquid flavours and continued vaping and tobacco smoking uptake or cessation.²⁵ The study found no evidence that flavoured e-liquid use versus unflavoured use was specifically associated with tobacco




Figure 2 Forest plot for e-cigarette use and smoking initiation in previous meta-analyses.*denotes risk ratios, otherwise odds ratios. n = number of studies pooled within each review. NR = not reported.

smoking uptake,²⁵ but with only three included studies, the authors recommended more research.

Smoking relapse

One systematic review studied relapse (the probability of transitioning from former to current cigarette smoking) and included young people, but only one longitudinal study (with a Markov model) specifically included young people.²¹ This primary study found that the probability of relapse was higher among young people using e-cigarettes in the past 30 days than those who had never used them. An umbrella review⁹ also focused on relapse, but included only adults.

Quality assessment

The quality of the included reviews ranged from high (one systematic review)²² or moderate (one systematic review⁴¹ and one umbrella review)⁸ to low (eight reviews) or critically low (13 reviews) (online supplemental table). The results were highly consistent, irrespective of quality assessment. Some reviews had fewer included studies, perhaps due to the date the searches were conducted or narrower inclusion criteria, such as limited populations studied (eg, Asian Americans³⁷ or Asian and Pacific Islanders),³⁶ a tight scope on flavoured vs unflavoured e-cigarettes,²⁵ or a sole focus on smoking relapse.²¹ However, many other reasons for the exclusion of primary studies were largely unexplained.

E-cigarette use and subsequent substance use

One umbrella review⁷ assessed associations between e-cigarette use and substance use. The review concluded that moderate evidence exists for associations between e-cigarette use and alcohol (OR 3.72, 95% CI 2.03 to 6.83) and marijuana use (OR 2.89, 95% CI 1.61 to 5.19). We included an additional systematic review⁴² to those included in the umbrella review⁷ and excluded one systematic review⁴³ in the umbrella review⁷ because it only

assessed how commonly e-cigarette technology was used as an illicit drug delivery system.

Five systematic reviews comprising 91 primary studies (61 unique) assessed the association between e-cigarette use and substance use; again, bidirectional relationships cannot be fully parsed, but these were mainly longitudinal studies with 1–7 year follow-up and cross-sectional surveys. Thirty-four per cent (21/61) of primary studies were included in multiple reviews (online supplemental table). The degree of overlap was high (CCA=12%). Country settings for primary studies were the US, UK, Australia, Sweden, Hong Kong, Poland, Russia, Iceland, Canada, Switzerland, China, Greece, France, Mexico, South Korea, Iceland, Taiwan and Germany. The total number of participants, where reported, ranged from 7551 to 373 150.

Marijuana use

Four systematic reviews demonstrated significantly greater odds of past or current e-cigarette use and subsequent marijuana use in adolescents and young adults, with adjusted OR 2.72, 95% CI 2.06 to 3.61⁴⁴, adjusted OR 3.47, 95% CI 2.63 to 4.59⁴⁵, OR 6.04, 95% CI 3.80 to 9.60⁴⁶ and OR 5.15, 95% CI 2.94 to 9.00,⁴² respectively (figure 3). This association was reported to be greater in younger populations in two reviews.^{45 46}

Alcohol use

Three systematic reviews identified significantly greater odds of alcohol use for young people using e-cigarettes compared with those never using them: OR 4.50, 95% CI 3.31 to 6.13⁴⁶, OR 6.62, 95% CI 5.67 to 7.72⁴⁷ and OR 6.67, 95% CI 2.85 to 15.59⁴² (online supplemental table). This relationship was also apparent with binge drinking: OR 4.51, 95% CI 3.13 to 6.51⁴⁶ and OR 6.73, 95% CI 4.50 to 10.07⁴⁷ (figure 3). As with cannabis use, the relationship between e-cigarette use and alcohol use was significantly greater in adolescent samples than adult samples (log OR 0.96 (OR 2.61), $z=4.21$, $p<0.001$).⁴⁶




Figure 3 Forest plot for all drug initiation use and abuse, and e-cigarette use versus never use. NR = adjusted or unadjusted status of effect size was not reported.

Stimulant use

One systematic review showed increased odds of non-prescribed Ritalin/Adderall use (OR 2.13, 95% CI 1.10 to 4.11) from one longitudinal cohort, but not for painkillers, sedatives or tranquilisers in young people using e-cigarettes compared with those never using them.⁴²

Quality assessment

All included reviews on substance use were rated as critically low quality due to significant methodological limitations. Issues included: lack of a priori methods, not listing excluded studies and inadequate reporting on risk of bias impacts. The searches in Lau *et al*⁴² were deemed problematic, which may help explain the small number of included studies in this, the most recent review identified.

Respiratory outcomes

One umbrella review⁷ reported on the impacts on respiratory outcomes (online supplemental table). This review was not limited to young people and included fewer systematic reviews related to young people than our umbrella review: two on asthma,^{48 49} two on e-cigarette or vaping product use associated lung injury (EVALI)^{50 51} and one on acute eosinophilic pneumonia.⁵² This umbrella review's pooled RR was 1.13 (95% CI 1.09 to 1.18), with the authors concluding that high level evidence exists that electronic nicotine delivery system use can exacerbate asthma.

Fifteen systematic reviews^{34 48–60} reported on the impacts on respiratory outcomes (online supplemental table). Primary studies were conducted in the US, South Korea, Canada and Kuwait. The total number of included participants, where reported, ranged from 162 182 to 483 948.

Asthma

Within the seven systematic reviews with asthma as an outcome, 47 primary studies were included, of which 20 were unique. Most studies were cross-sectional and the degree of overlap was very high (CCA=23%).

In all meta-analyses conducted within the reviews, statistically significant increases in asthma diagnosis or exacerbation were observed with e-cigarette use (whether measured as any use, ever use or current use) (figure 4). Overall, the reviews concluded that there was an increased risk of young people who vape developing asthma or asthma exacerbation, with pooled adjusted ORs of 1.20–1.36 for diagnosis and 1.44 for exacerbation.

Respiratory outcomes (excluding asthma)

One systematic review of the effect of e-cigarette use on coughing included six cross-sectional surveys, and one retrospective medical chart review from the US, Canada, Switzerland and Hong Kong, which found that adolescent use of e-cigarettes was associated with increased coughing, and young people who vape are more likely to report coughing compared with those who do not vape.⁵³ There was no pooled meta-analysis, but higher ORs were reported for young people with more frequent use (adjusted OR range 1.30–2.13). Four of these cross-sectional surveys were reported in another review.⁶⁰ There was no pooled meta-analysis, but the authors concluded that there was an association between e-cigarette use and effects on respiratory systems.

Three reviews^{48 50 51} reported on case series of EVALI of upwards of 53 cases, showing that these cases had occurred in patients as young as 13 years,⁵¹ with a wide range of outcomes,




Figure 4 Forest plot for meta-analyses of e-cigarette use and asthma diagnosis or exacerbation versus never use. Ever use, current use and use reflect terminology used in original reviews. *Chand *et al*: ever use of e-cigarettes and asthma diagnosis - includes two studies on adults (18+); current use of e-cigarettes and asthma - includes one study on adults (18+).

and that e-cigarette use was "involved in damage related to the development of pulmonary disease", with e-liquids containing tetrahydrocannabinol (THC) a feature in 80% of cases.⁵⁰

Other systematic reviews reported cross-sectional data on increased wheezing,⁶⁰ dyspnoea⁶⁰ and bronchitis,^{58 60} and potential increases in allergic rhinitis.⁶⁰ Four reviews^{34 48 52 59} highlighted respiratory problems (pneumonia, bronchitis and pneumothorax) using case reports.

Quality assessment

All respiratory outcome reviews were rated as critically low quality, except two on asthma which were rated as low quality.^{54 57} The lack of an adequate search strategy in the reviews may help explain the low number of included studies, while lack of overlap was partly explained by different exposure or outcome measures.

Burns and other injuries

One umbrella review⁶ examined risk of injuries resulting from using e-cigarettes and reported on five US case reports in young people (four explosions and one inadvertent aspiration). Another umbrella review⁷ included three systematic reviews^{48 61 62} that we included but reported no young person-specific results or conclusions.

Six systematic reviews^{48 59 62–65} including 34 individual studies examined the risk of injuries resulting from using e-cigarettes in young people. Primary studies were conducted in the US, UK, France, Canada, Germany and Malaysia. The degree of overlap was very high (CCA=24%).

The six reviews predominantly included case reports and case series, as well as three surveillance system based studies. The reviews reported incidents of explosions among young people, resulting in projectile and burn injuries. Body parts affected were the mouth/face and hands, when e-cigarettes exploded mid-use, as well as lower limbs, when e-cigarettes exploded in pockets. Other less frequent injuries involved inadvertent aspiration of e-cigarette parts (outcomes reported alongside these injuries related to accidental or intentional poisoning were excluded

given our review's focus on e-cigarette devices used as intended by manufacturers).

Quality assessment

Five of the included systematic reviews were rated as critically low quality, while the remaining systematic review and the one umbrella review were rated as low quality. The most common weaknesses were not using a PECO framework, not addressing heterogeneity (perhaps reflecting the lack of statistical analyses due to case report data) and not providing lists of excluded studies.

Mental health

Two umbrella reviews^{7 8} included one of the systematic reviews that we included (Becker *et al*⁶⁶ in both cases), with one⁸ combining these findings with that of an individual study.⁴⁵ A further umbrella review⁶ included a study (Marsden *et al*) found in our other systematic reviews, but reported no young person specific or mental health related results.⁶

Seven systematic reviews^{57 58 66–70} included 63 individual studies (of which 54 were unique) examining associations between vaping and mental health. Studies were predominantly cross-sectional surveys, which complicates assessments of directionality/causality, and longitudinal studies. There was little overlap between these reviews (CCA=4%). Primary studies were conducted in the US, South Korea, UK and Taiwan.

Depression and suicidal outcomes

One umbrella review found,⁷ as per the Becker *et al* review that it was based on,⁶⁶ that e-cigarette use was associated with greater depression and suicidality (as well as attention deficit/hyperactivity disorder, eating disorders and stress) among adolescents. The other umbrella review⁸ was similarly focused predominantly on Becker *et al*, concluding that e-cigarette-only use compared with no use was associated with depressive symptoms (adjusted OR 1.37, 95% CI 1.19 to 1.57) and that past 30-day e-cigarette use was associated with suicide ideation (adjusted OR 2.49, 95% CI 1.82 to 3.42), suicide planning (adjusted OR 4.63, 95% CI 3.22 to 6.67) and suicide attempts (adjusted OR 6.17, 95% CI 4.13 to 9.24), with ORs for suicide plans greater in female adolescents.

The only systematic review to incorporate a meta-analysis⁶⁷ found a pooled OR for children and adolescents who vape of 1.46 (95% CI 1.33 to 1.59) for suicide ideation, 2.23 (95% CI 1.74 to 2.71) for suicide planning and 2.46 (95% CI 1.94 to 2.98) for suicide attempts. One further review found that e-cigarette use was associated with depression and suicidality in adolescents,⁵⁸ while another⁶⁸ found conflicting results in relation to depression associations in this age group from its included studies.

Other mental health related outcomes

Other systematic reviews synthesised evidence on adolescent weight concerns⁶⁹ - finding that higher concerns were associated with higher frequency of e-cigarette use - or sleep complaints.^{57 70} The latter two reviews found respectively that adolescents who vaped had shorter total sleep time on weekends and that those who used both e-cigarettes and conventional cigarettes reported insufficient sleep significantly more often than those who used conventional cigarettes only.

Given the heterogeneity in outcomes involved, a meta-analysis was not possible; the absence of meta-analyses in all but one systematic review also precluded a forest plot for this outcome.

Quality assessment

One included umbrella review was rated as moderate quality, while the other was critically low. Of the seven systematic reviews, two were rated as low quality while the others were critically low. Common weaknesses were not addressing heterogeneity, not reporting included studies' funding sources and not justifying study design.

Cardiovascular disorders

Three systematic reviews included studies examining associations between use of e-cigarettes and cardiovascular outcomes in youth.^{48 59 65} One umbrella review⁷ included a section on cardiovascular disease, but its included systematic reviews were all excluded from ours because none undertook young person specific analyses, while its own cardiovascular section similarly included no young person specific results.

The three systematic reviews only incorporated two individual studies reporting cardiovascular outcomes in young people: a case report from Turkey⁷¹ and a cross-sectional study of 270 Russian students.⁷² The cross-sectional study found that an arterial stiffness marker was significantly higher in young people who vaped than those who did not. The case report dated from 2014 and involved acute myocardial ischaemia in a 24-year-old man with an absence of other risk factors; the report concluded that nicotine in the e-liquid may have been the cause.

Quality assessment

Of the three included systematic reviews, one was rated low quality and the other two were critically low. As elsewhere, common issues were not listing excluded studies and not adequately considering heterogeneity.

Oral health

Livingston *et al*⁵⁸ and Bozier *et al*³⁴ both included a cross-sectional study⁷³ of 65 528 students in South Korea on oral health that indicated that e-cigarette use was associated with significantly increased odds of gingival pain and/or bleeding, tongue and/or inside cheek pain and cracked or broken teeth. The review by Livingston *et al* also reported on the PATH longitudinal study⁷⁴ which indicated that, among 13 650 participants aged 12–17 years in the US, e-cigarette use alone was not significantly more likely than non-use to be associated with self-reported diagnosed dental problems. These reviews did not report comprehensive search strategies.

Guo *et al*⁷⁵ included six cross-sectional studies and one case-control study with 6827 young participants and found an association with e-cigarette use and dry mouth (xerostomia), with a higher prevalence in those aged <25 years than those aged >25 years (30%, 95% CI 28 to 32 vs 20%, 95% CI 18 to 23). One of these cross-sectional surveys⁷⁶ (975 US adolescents aged 13–17 years) was also reported by Kurdys-Bykowska *et al*⁶⁰ alongside one of 3488 children (aged 6–12 years) in Zurich,⁷⁷ which also reported higher rates of dry mouth. Another review reported a higher risk of throat dryness in young people who vaped (from one cross-sectional survey⁷⁸ of 520 paediatricians collecting information on children and adolescents in Canada).⁶¹

Other harm outcomes

Two systematic reviews^{65 79} included the same cross-sectional study⁸⁰ of 1221 Danish young men that found that those who used e-cigarettes daily had significantly lower total sperm counts than those who did not use e-cigarettes. This study was rated a low risk of bias by the review authors.⁶⁵

One review reporting the results of a cross-sectional survey of 975 US adolescents⁷⁶ indicated that dizziness or lightheadedness and headaches or migraines were more likely to be reported among those using e-cigarettes compared with those not using them (43.8% vs 14.6%, $p=0.002$).⁶⁰

Valadez-Cuen *et al*⁸¹ concluded that people using e-cigarettes were at a higher risk of contracting COVID-19, but only one cross-sectional online survey of 4351 participants was restricted to young people aged 13–24 years.⁸² This review was rated as critically low quality and did not report a comprehensive search strategy.

There was limited evidence for other harms. One systematic review⁷⁹ reported on a cross-sectional study⁸³ which concluded that, among 441 adolescents and young adults with paediatric onset chronic kidney disease (CKD), occasional e-cigarette use was not associated with disease progression, proteinuria or elevated blood pressure. In the same review, another cross-sectional study⁷² found e-cigarette use was related to higher albuminuria values (an early marker of CKD) after measuring urine albumin levels in 270 young volunteers. One systematic review⁸⁴ assessed the implications of vaping on surgical wound healing; one of the case reports⁸⁵ included was of a patient aged <25 years with acute e-cigarette consumption affecting flap tissue oximetry due to acute vasoconstriction.

DISCUSSION

The systematic reviews included in this umbrella review, which have primarily examined longitudinal, individual level studies, consistently indicated a significant association between e-cigarette use and later cigarette smoking in young people. These repeated strong associations in prospective cohort studies are consistent with a causal relationship between vaping and subsequent smoking. Inferring causality remains challenging,²⁶ and authors' conclusions within the reviews vary as to whether the relationship between e-cigarettes and smoking is causal,^{20 27} reflecting the contested/uncertain nature of this interpretation elsewhere and plausible competing explanations for these associations. Nonetheless, given the consistent associations we observed with increased smoking and multiple possible harms to health and wellbeing in this age group, which are consistent with possible causal effects, the evidence supports policy measures to protect young people who do not smoke from the potential risks associated with vaping. These are therefore proposed on a precautionary basis in light of definitive evidence of causality not yet being established.

Multiple theories attempt to explain the relationship between e-cigarette use and smoking: the gateway effect⁸⁶ (vaping may increase subsequent smoking risk), the diversion effect⁸⁷ (vaping may reduce smoking by creating an alternative) and the common liability hypothesis⁸⁸ (no causal relationship). The consistency in the evidence we present is striking; e-cigarette use raises the likelihood of future cigarette smoking among youth and young adults. It may also increase smoking frequency and intensity (ie, the gateway effect). While population data show opposing trends, with smoking rates generally falling as vaping rises, theoretically supporting a diversion effect,^{23 88 89} smoking trends measured in population studies are also influenced by other factors, such as public smoking bans, taxation and restrictions on purchasing and packaging. Our findings also provide evidence against the diversion effect because, under this hypothesis, people vaping would be less likely to smoke than those not vaping.

We also found associations between e-cigarette use and subsequent substance use. Certain behaviours, such as e-cigarette,

alcohol and marijuana use, often cluster, potentially fulfilling similar social and psychological roles.^{24 90 91} Many of the included studies within the reviews used adjusted ORs where available, which sometimes extended beyond assessing sex, gender and socioeconomic status variables. Many studies also used longitudinal designs with long follow-up periods, suggesting a gateway theory as opposed to a reverse gateway theory. This is supported by one of the systematic reviews⁴³ included in Asfar *et al*⁷ which indicated that vape equipment can be used as a drug delivery system for illicit drugs and found that, of 861 respondents, 39.5% of people using electronic nicotine delivery systems had used them to vape recreational drugs, with marijuana being the most common (18.0%).

Our findings furthermore suggest that there is a link between current and ever use of e-cigarettes and an asthma diagnosis or exacerbation of existing asthma. In studies investigating e-cigarettes in real world settings, it is difficult to eliminate confounding due to the complex nature of these settings and the limitations of observational studies. However, the consistency in the findings indicates at least an association between e-cigarettes and increased asthma risk.

Acute exposure to e-cigarette vapour can cause coughing, bronchospasm and inflammatory responses in airways.^{92 93} However, evidence for other respiratory harms associated with e-cigarette use in young people is lacking. While the evidence we present suggests that coughing may be associated with e-cigarette use,⁵³ more research is required for pneumonia,⁵² bronchitis⁵⁸ and allergic rhinitis. Case reports and case series were identified for EVALI within the reviews. Most of these cases have been connected with the use of e-liquids containing THC and vitamin E acetate. However, many more case reports have been reported to regulatory agencies in the US and elsewhere, not all implicating THC.^{94–96}

Our review also found repeated significant associations between e-cigarette use and mental health outcomes. The two umbrella reviews to date have largely been based on a single systematic review (Becker *et al*),⁶⁶ and those authors rate their evidence for links with depression as weak to moderate and for links with suicide as weak, citing the frequency of cross-sectional designs and possible confounding. Given the severity of these outcomes, urgent longitudinal research is merited. Other areas for further research include oral health, fertility, headaches or migraines, and dizziness or lightheadedness, while the limited number of reviews incorporating cardiovascular outcomes likely reflects our focus on youth populations where cardiovascular events are rarer than in adults.

A serious common harm of e-cigarettes is addiction. However, addiction as an outcome was noticeably absent from included reviews. This may be because it is an established property of nicotine and thus deemed self-evident. However, nicotine is one of the most addictive substances known, and there is clear evidence that young people are experiencing addiction to it as a direct consequence of use of e-cigarettes, including having difficulty in quitting.^{97–99} Being caught in the addiction cycle of withdrawal, craving and satiation can significantly impact daily functioning¹⁰⁰ and children's behaviour in schools, as well as increasing social exclusion and stigma.^{101 102} These areas of concern remain important avenues for future research.

Another striking gap in the evidence we collected, given the considerable interest in the role of nicotine on neurodevelopment,^{103 104} is the potential influence of e-cigarettes on adolescent brain development. Only one umbrella review considered this in relation to young people (Banks *et al*)⁶ but found no evidence concerning e-cigarettes and brain development (or neurological

outcomes generally). Systematic reviews assessing tobacco use and neurocognitive impairment^{105 106} indicate that findings from research on young people may differ from adults. A previous non-systematic review synthesised the evidence on e-cigarettes and brain development, suggesting harmful effects,¹⁰⁷ and a scoping review found that e-cigarette use among young people is associated with poor academic achievement, with a lack of evidence on the causal direction.¹⁰⁸

Use of both e-cigarettes and cigarettes together is also increasingly prevalent and was not the focus of this umbrella review. However, this dual use delivers a wider range of toxins than solely smoking or vaping; evidence from a systematic review indicates that risks of harm may be higher than using either product alone.¹⁰⁹

Most of the systematic reviews we included were rated as low or critically low quality using AMSTAR 2 (multiple analyses have found that AMSTAR 2 rates most reviews as critically low).^{110–112} As with all quality assessments, it was challenging to distinguish the quality of systematic review reporting from the quality of the review's conduct.¹¹¹ An issue for most reviews was the lack of an excluded studies list (with reasons for exclusion). This is a critical item in AMSTAR 2 and therefore resulted in downgrading of many of the reviews.

Strengths and weaknesses

While we were able to identify other umbrella reviews on the topic of harmful effects of e-cigarettes, none focused on a full range of potential harms in adolescents or young people. Indeed, our umbrella review is, to our knowledge, the most comprehensive synthesis of systematic reviews that has taken place to date on the wider risks of vaping in young people.

Another strength of our research is that, once early findings began to emerge, we discussed them with young people directly at a youth public health forum. Participants advised on how their age group might view our results, citing frustration at instances elsewhere when vaping is portrayed as much safer than smoking, and how we could best disseminate results effectively.

As with all umbrella reviews, we were dependent on the quality of the included systematic reviews, and our results lack some of the granular detail that can be ascertained in a single systematic review. Much of the evidence available for outcomes other than smoking uptake is cross-sectional; there was a paucity of population based (ecological) analyses which could have helped assess if changes in vaping prevalence and smoking prevalence were associated at the population level. Furthermore, although we restricted our searches to 2015 onwards (see methods), there were some limited data within the included reviews that predated 2015. We have, however, provided a comprehensive high-level overview of the research on harmful impacts of e-cigarettes in young people and, by measuring the overlap in included studies, have been able to provide an overview for the risk of inflating research results when primary studies' findings are shared multiple times.

Future evidence

There were 12 systematic reviews registered on PROSPERO between 2020 and 2024 categorised as ongoing and aiming to evaluate the harms of e-cigarettes in young people (although many reviews registered on PROSPERO might never be completed or published). Three focus on respiratory outcomes^{113–115} and one on asthma.¹¹⁶ One review is examining smoking initiation¹¹⁷ but, given our consistent findings on this harm, it is unlikely to change the conclusions of our review. Two focus on substance

use,^{118 119} one on dental caries,¹²⁰ one on depression¹²¹ and one on physical activity.¹²² Two reviews are investigating general harms.^{123 124}

Policy implications

Our findings support the implementation of policy measures to restrict sales and marketing of e-cigarettes to young people, and restrictions on advertising design features that are designed for, or likely to appeal to, young people. Such efforts may form part of a wider set of measures to restrict harms, including raising the public's and young people's awareness of these harms, and counter-marketing to raise public and policy awareness of the marketing and strategies that e-cigarette companies have targeted at children and young people. Little research focuses on evidence regarding access to e-cigarettes. In an evidence gap map,¹²⁵ seven studies were identified on taxation and other price policies, and two on clean indoor air policies.

CONCLUSIONS

Systematic reviews showed consistent evidence that higher risks of smoking initiation, substance use (marijuana, alcohol and stimulants), asthma, coughing, injuries and adverse mental health outcomes are associated with e-cigarette use in young people. Other harms such as pneumonia, bronchitis, headaches, migraines, dizziness/lightheadedness, low sperm count and oral health harms are possibly associated with e-cigarette use in young people but require further research.

Acknowledgements We thank the Youth Advisory Panel of the Association for Young People's Health (AYPH) for their contributions to patient and public involvement discussions on this work, and Edwin van Leeuwen for his R code.

Contributors SG and GH are joint first authors. SG ran searches, conducted study selection, data extraction, critical appraisal, secondary checking of others' data extraction and critical appraisal, and analysis, and wrote the manuscript. GH conducted study selection, data extraction, critical appraisal, secondary checking of others' data extraction and critical appraisal, and analysis, and wrote the manuscript. LMB and SGN conducted data extraction, critical appraisal, secondary checking of others' data extraction and critical appraisal, and analysis. REG designed the study, conducted study selection and analysis, and supervised the project. MP designed the study and supervised the project. All authors critically reviewed and edited the manuscript, approved the final version for submission, and agreed to be accountable for all aspects of this work. REG is the guarantor. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

Funding The study is independent research carried out by the Public Health Policy Research Unit (PH-PRU), grant No PH_PRU_1217_20901, commissioned and funded by the National Institute for Health and Social Care Research (NIHR) Policy Research Programme. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care (DHSC). MP is co-investigator in the SPECTRUM consortium which is funded by the UK Prevention Research Partnership (UKPRP), a consortium of UK funders (UKRI Research Councils: Medical Research Council (MRC), Engineering and Physical Sciences Research Council (EPSRC), Economic and Social Research Council (ESRC) and Natural Environment Research Council (NERC); charities: British Heart Foundation, Cancer Research UK, Wellcome and the Health Foundation; government: Scottish Government Chief Scientist Office, Health and Care Research Wales, NIHR and Public Health Agency (NI)). SG is employed by the University of York and has worked on grants from NIH, NIHR, DHSC, MRC and Wellcome. SG has authored Cochrane Reviews and a Cochrane handbook. GH has received funding from NIHR and ESRC and, as a clinical academic, holds an honorary contract with the Office for Health Improvement and Disparities (OHID). REG has accepted funding from NIHR and MRC and accepted honoraria from WHO and the Academy of Medical Sciences. LMB and SGN are Public Health Specialty Registrars and writing in their capacity as Honorary Research Fellows at the London School of Hygiene and Tropical Medicine. SGN also has an honorary contract with the UK Health Security Agency (UKHSA).

Competing interests All authors have completed the ICMJE uniform disclosure form at www.icmje.org/disclosure-of-interest/ and declare: funding for the study as detailed above; no financial relationships with any organisations that might have an

interest in the submitted work in the previous three years; no other relationships or activities that could have influenced the submitted work.

Patient consent for publication Not applicable.

Ethics approval As this umbrella review collated and analysed data from publicly accessible materials, this study did not require ethical approval.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All the systematic reviews and umbrella reviews included in this review are published. The complete data extraction sheets are available in supplementary files.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Greg Hartwell <http://orcid.org/0000-0001-9153-7126>

Rebecca E Glover <http://orcid.org/0000-0001-9150-9977>

REFERENCES

- 1 CDC. Smoking and tobacco use: e-cigarette use among youth: U.S. Centers for Disease Control and Prevention; 2024. Available: <https://www.cdc.gov/tobacco/e-cigarettes/youth.html> [Accessed 4 Jun 2025].
- 2 Tobacco: e-cigarettes. World Health Organization; 2024. Available: <https://www.who.int/news-room/questions-and-answers/item/tobacco-e-cigarettes>
- 3 World Health Organization. Prevalence of tobacco and e-cigarette use by young people in the WHO European region in 2022. WHO; 2024. Available: <https://www.who.int/europe/publications/m/item/prevalence-of-tobacco-and-e-cigarette-use-by-young-people-2022> [Accessed 4 Jun 2025].
- 4 NHS. Vaping to quit smoking. Better Health: Let's do this; 2024. Available: <https://www.nhs.uk/better-health/quit-smoking/ready-to-quit-smoking/vaping-to-quit-smoking>
- 5 Mahase E. MPs back bill that will phase out smoking and restrict vaping. *BMJ* 2024;387:q2668.
- 6 Banks E, Yazdijoglou A, Brown S, et al. Electronic cigarettes and health outcomes: umbrella and systematic review of the global evidence. *Med J Aust* 2023;218:267–75.
- 7 Asfar T, Jebai R, Li W, et al. Risk and safety profile of electronic nicotine delivery systems (ENDS): an umbrella review to inform ENDS health communication strategies. *Tob Control* 2022.
- 8 Khan AM, Ahmed S, Sarfraz Z, et al. Vaping and Mental Health Conditions in Children: An Umbrella Review. *Subst Abuse* 2023;17.
- 9 Baenziger ON, Ford L, Yazdijoglou A, et al. E-cigarette use and combustible tobacco cigarette smoking uptake among non-smokers, including relapse in former smokers: umbrella review, systematic review and meta-analysis. *BMJ Open* 2021;11:e045603.
- 10 Gates M, Gates A, Pieper D, et al. Reporting guideline for overviews of reviews of healthcare interventions: development of the PRIOR statement. *BMJ* 2022;378:e070849.
- 11 KSR evidence: searching. KSR Evidence; 2024. Available: <https://ksrevidence.com/>
- 12 Use of vapes (e-cigarettes) among young people in Great Britain. Action on Smoking and Health (ASH); 2024. Available: <https://ash.org.uk/uploads/Use-of-vapes-among-young-people-in-Great-Britain-2024.pdf?v=1738083753>
- 13 EndNote. Clarivate; 2024. Available: <https://endnote.com>
- 14 Covidence systematic review software. Melbourne, Australia Veritas Health Innovation; 2024. Available: <https://www.covidence.org>
- 15 Shea BJ, Reeves BC, Wells G, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ* 2017;358:j4008.
- 16 Hennessy EA, Johnson BT. Examining overlap of included studies in meta-reviews: Guidance for using the corrected covered area index. *Res Synth Methods* 2020;11:134–45.
- 17 Aladeokin A, Haighton C. Is adolescent e-cigarette use associated with smoking in the United Kingdom?: A systematic review with meta-analysis. *Tob Prev Cessat* 2019;5:15.
- 18 Chan GCK, Stjepanović D, Lim C, et al. Gateway or common liability? A systematic review and meta-analysis of studies of adolescent e-cigarette use and future smoking initiation. *Addiction* 2021;116:743–56.
- 19 Chatterjee K, Alzghoul B, Innabi A, et al. Is vaping a gateway to smoking: a review of the longitudinal studies. *Int J Adolesc Med Health* 2016;30.
- 20 Khouja JN, Suddell SF, Peters SE, et al. Is e-cigarette use in non-smoking young adults associated with later smoking? A systematic review and meta-analysis. *Tob Control* 2021;30:8–15.
- 21 Kim MM, Steffensen I, Miguel RTD, et al. A Systematic Review Investigating Associations Between E-Cigarette Use Among Former Cigarette Smokers and Relapse to Smoking Cigarettes. *Inquiry* 2023;60:469580231214457.
- 22 Kim MM, Steffensen I, Miguel RTD, et al. A systematic review and meta-analysis of the association between e-cigarette use among non-tobacco users and initiating smoking of combustible cigarettes. *Harm Reduct J* 2024;21.
- 23 Levy DT, Warner KE, Cummings KM, et al. Examining the relationship of vaping to smoking initiation among US youth and young adults: a reality check. *Tob Control* 2019;28:629:629–35.
- 24 McCabe SE, West BT, Veliz P, et al. E-cigarette Use, Cigarette Smoking, Dual Use, and Problem Behaviors Among U.S. Adolescents: Results From a National Survey. *J Adolesc Health* 2017;61:155–62.
- 25 Notley C, Gentry S, Cox S, et al. Youth use of e-liquid flavours—a systematic review exploring patterns of use of e-liquid flavours and associations with continued vaping, tobacco smoking uptake or cessation. *Addiction* 2022;117:1258–72.
- 26 O'Brien D, Long J, Quigley J, et al. Association between electronic cigarette use and tobacco cigarette smoking initiation in adolescents: a systematic review and meta-analysis. *BMC Public Health* 2021;21:954.
- 27 Soneji S, Barrington-Trimis JL, Wills TA, et al. Association Between Initial Use of e-Cigarettes and Subsequent Cigarette Smoking Among Adolescents and Young Adults: A Systematic Review and Meta-analysis. *JAMA Pediatr* 2017;171:788–97.
- 28 Associations between the use of snus (moist tobacco) or electronic cigarettes and tobacco smoking: a systematic review. Swedish Agency for Health Technology A, Assessment of Social S, 2020:312.
- 29 Zhang Y-Y, Bu F-L, Dong F, et al. The effect of e-cigarettes on smoking cessation and cigarette smoking initiation: An evidence-based rapid review and meta-analysis. *Tob Induc Dis* 2021;19:04.
- 30 Dautzenberg B, Legleye S, Underner M, et al. Systematic Review and Critical Analysis of Longitudinal Studies Assessing Effect of E-Cigarettes on Cigarette Initiation among Adolescent Never-Smokers. *Int J Environ Res Public Health* 2023;20:6936.
- 31 Yoong SL, Hall A, Turon H, et al. Association between electronic nicotine delivery systems and electronic non-nicotine delivery systems with initiation of tobacco use in individuals aged < 20 years. A systematic review and meta-analysis. *PLoS ONE* 2021;16.
- 32 Adermark L, Galanti MR, Ryk C, et al. Prospective association between use of electronic cigarettes and use of conventional cigarettes: a systematic review and meta-analysis. *ERJ Open Res* 2021;7:00976–2020.
- 33 Barufaldi LA, Guerra RL, Albuquerque R de C de, et al. Risk of initiation to smoking with the use of electronic cigarettes: systematic review and meta-analysis. *Cien Saude Colet* 2021;26:6089–103.
- 34 Bozler J, Chivers EK, Chapman DG, et al. The Evolving Landscape of e-Cigarettes: A Systematic Review of Recent Evidence. *Chest* 2020;157:1362–90.
- 35 Health I, Quality A. Health technology assessment (HTA) of smoking cessation interventions. 2017;391.
- 36 Phillips KT, Okamoto SK, Johnson DL, et al. Correlates of tobacco use among Asian and Pacific Islander youth and young adults in the U.S.: A systematic review of the literature. *Exp Clin Psychopharmacol* 2021;29:440–55.
- 37 Shi M, Gette JA, Gissandner TD, et al. E-cigarette use among Asian Americans: a systematic review. *J Ethn Subst Abuse* 2020;ahead of print:1–34.
- 38 Zhong J, Cao S, Gong W, et al. Electronic Cigarettes Use and Intention to Cigarette Smoking among Never-Smoking Adolescents and Young Adults: A Meta-Analysis. *Int J Environ Res Public Health* 2016;13:465.
- 39 Pieper D, Antoine S-L, Mathes T, et al. Systematic review finds overlapping reviews were not mentioned in every other overview. *J Clin Epidemiol* 2014;67:368–75.
- 40 Pepper JK, Ribisl KM, Brewer NT. Adolescents' interest in trying flavoured e-cigarettes. *Tob Control* 2016;25:ii62–6.
- 41 NICE. Evidence Reviews for E-Cigarettes and Young People: Tobacco: Preventing Uptake, Promoting Quitting and Treating Dependence: Update: Evidence Review F and G. 2021. Available: <https://www.nice.org.uk/guidance/ng209/evidence/f-g-e-cigarettes-and-young-people-pdf-10890777857>
- 42 Lau L, Conti AA, Hemmati Z, et al. The prospective association between the use of E-cigarettes and other psychoactive substances in young people: A systematic review and meta-analysis. *Neurosci Biobehav Rev* 2023;153:105392.
- 43 Breitbarth AK, Morgan J, Jones AL. E-cigarettes—An unintended illicit drug delivery system. *Drug Alcohol Depend* 2018;192:98–111.
- 44 Yimer TM, McClure-Thomas C, Stjepanovic D, et al. The relationship between cannabis and nicotine use: A systematic review and meta-analysis. *Addiction* 2024;119:2076–87.

- 45 Chadi N, Schroeder R, Jensen JW, *et al.* Association Between Electronic Cigarette Use and Marijuana Use Among Adolescents and Young Adults: A Systematic Review and Meta-analysis. *JAMA Pediatr* 2019;173:e192574.
- 46 Hershberger A, Argyriou E, Cyders M. Electronic nicotine delivery system use is related to higher odds of alcohol and marijuana use in adolescents: Meta-analytic evidence. *Addict Behav* 2020;105:106325.
- 47 Rothrock AN, Andris H, Swetland SB, *et al.* Association of E-cigarettes with adolescent alcohol use and binge drinking-drunkness: A systematic review and meta-analysis. *Am J Drug Alcohol Abuse* 2020;46:684–98.
- 48 Tzortzi A, Kapetanstratiki M, Evangelopoulou V, *et al.* A Systematic Literature Review of E-Cigarette-Related Illness and Injury: Not Just for the Respiriologist. *Int J Environ Res Public Health* 2020;17:E22482248.
- 49 Xian S, Chen Y. E-cigarette users are associated with asthma disease: A meta-analysis. *Clin Respir J* 2021;15:457–66.
- 50 Bravo-Gutiérrez OA, Falfán-Valencia R, Ramírez-Venegas A, *et al.* Lung Damage Caused by Heated Tobacco Products and Electronic Nicotine Delivery Systems: A Systematic Review. *Int J Environ Res Public Health* 2021;18:4079.
- 51 Gonsalves CL, Zhu JW, Kam AJ. Diagnosis and Acute Management of E-Cigarette or Vaping Product Use—Associated Lung Injury in the Pediatric Population: A Systematic Review. *J Pediatr* 2021;228:260–70.
- 52 Chaaban T. Acute eosinophilic pneumonia associated with non-cigarette smoking products: a systematic review. *Adv Respir Med* 2020;88:142–6.
- 53 Bourke M, Sharif N, Narayan O. Association between electronic cigarette use in children and adolescents and coughing a systematic review. *Pediatr Pulmonol* 2021;56:3402–9.
- 54 Chand BR, Hosseinzadeh H. Association between e-cigarette use and asthma: a systematic review and meta-analysis. *J Asthma* 2022;59:1722–31.
- 55 Li X, Zhang Y, Zhang R, *et al.* Association between E-Cigarettes and Asthma in Adolescents: A Systematic Review and Meta-Analysis. *Am J Prev Med* 2022;62:953–60.
- 56 Rocha A, Miyawaki AE, Trombini MA, *et al.* Risk of asthma exacerbation in adolescent users of electronic nicotine delivery systems: a systematic review and meta-analysis. *Arq Asma Alerg Immunol* 2023;7:41–8.
- 57 Pisinger C, Rasmussen SKB. The Health Effects of Real-World Dual Use of Electronic and Conventional Cigarettes versus the Health Effects of Exclusive Smoking of Conventional Cigarettes: A Systematic Review. *Int J Environ Res Public Health* 2022;19:13687.
- 58 Livingston JA, Chen CH, Kwon M, *et al.* Physical and mental health outcomes associated with adolescent E-cigarette use. *J Pediatr Nurs* 2022;64:1–17.
- 59 Hua M, Talbot P. Potential health effects of electronic cigarettes: A systematic review of case reports. *Prev Med Rep* 2016;4:169–78.
- 60 Kurdyś-Bukowska P, Koźmider L, Konwant D, *et al.* E-cigarettes use and respiratory symptoms in asthmatic and non-asthmatic adolescents – a systematic review. *Pediatr Med Rodz* 2024;20:29–36.
- 61 Yang I, Sandeep S, Rodriguez J. The oral health impact of electronic cigarette use: a systematic review. *Crit Rev Toxicol* 2020;50:97–127.
- 62 Seitz CM, Kabir Z. Burn injuries caused by e-cigarette explosions: A systematic review of published cases. *Tob Prev Cessat* 2018;4:32.
- 63 Jones CD, Ho W, Gunn E, *et al.* E-cigarette burn injuries: Comprehensive review and management guidelines proposal. *Burns* 2019;45:763–71.
- 64 Tran V, Mian M, Sreedharan S, *et al.* Oral and Maxillofacial Injuries Associated With E-Cigarette Explosions: A Systematic Review and Management Guidelines Proposal. *J Oral Maxillofac Surg* 2023;81:583–92.
- 65 McNeill A, Brose L, Robson D, *et al.* Nicotine vaping in England: an evidence update including health risks and perceptions. 2022. Available: <https://assets.publishing.service.gov.uk/media/633469f8fa8f5066d28e1a2/Nicotine-vaping-in-England-2022-report.pdf>
- 66 Becker TD, Arnold MK, Ro V, *et al.* Systematic Review of Electronic Cigarette Use (Vaping) and Mental Health Comorbidity Among Adolescents and Young Adults. *Nicotine Tob Res* 2021;23:415–25.
- 67 Awad AA, Itumalla R, Gaidhane AM, *et al.* Association of electronic cigarette use and suicidal behaviors: a systematic review and meta-analysis. *BMC Psychiatry* 2024;24:608.
- 68 Farooqui M, Shoaib S, Afaq H, *et al.* Bidirectionality of smoking and depression in adolescents: a systematic review. *Trends Psychiatry Psychother* 2023.
- 69 Hod R, Mohd Nor NH, Maniam S. Systematic review on e-cigarette and its effects on weight gain and adipocytes. *PLoS One* 2022;17:e0270818.
- 70 Kwon M, Park E, Dickerson SS. Adolescent substance use and its association to sleep disturbances: A systematic review. *Sleep Health* 2019;5:382–94.
- 71 Kivrak T, Sunbul M, Durmus E, *et al.* Acute myocardial infarction due to liquid nicotine in a young man. *Ther Adv Cardiovasc Dis* 2014;8:32–4.
- 72 Podzolkov VI, Bragina AE, Druzhinina NA, *et al.* Relation between Tobacco Smoking/ Electronic Smoking and Albuminuria/Vascular Stiffness in Young People without Cardiovascular Diseases. *Kidney Blood Press Res* 2020;45:467–76.
- 73 Cho JH. The association between electronic-cigarette use and self-reported oral symptoms including cracked or broken teeth and tongue and/or inside-cheek pain among adolescents: A cross-sectional study. *PLoS One* 2017;12:e0180506.
- 74 Akinkugbe AA. Cigarettes, E-cigarettes, and Adolescents' Oral Health: Findings from the Population Assessment of Tobacco and Health (PATH) Study. *JDR Clin Translational Res* 2019;4:276–83.
- 75 Guo X, Hou L, Peng X, *et al.* The prevalence of xerostomia among e-cigarette or combustible tobacco users: A systematic review and meta-analysis. *Tob Induc Dis* 2023;21:22.
- 76 King JL, Reboussin BA, Merten JW, *et al.* Negative health symptoms reported by youth e-cigarette users: Results from a national survey of US youth. *Addict Behav* 2020;104:106315.
- 77 Mozun R, Ardura-Garcia C, de Jong CCM, *et al.* Cigarette, shisha, and electronic smoking and respiratory symptoms in Swiss children: The LUIS study. *Pediatr Pulmonol* 2020;55:2806–15.
- 78 Richmond SA, Pike I, Maguire JL, *et al.* E-cigarettes: A new hazard for children and adolescents. *Paediatr Child Health* 2018;23:255–9.
- 79 Bandara NA, Zhou XR, Alhamam A, *et al.* The genitourinary impacts of electronic cigarette use: a systematic review of the literature. *World J Urol* 2023;41:2637–46.
- 80 Holmboe SA, Priskorn L, Jensen TK, *et al.* Use of e-cigarettes associated with lower sperm counts in a cross-sectional study of young men from the general population. *Hum Reprod* 2020;35:1693–701.
- 81 Valadez-Cuen K, Bhatt T, Mendez IE, *et al.* E-cigarette Use and Severe Coronavirus Disease 2019 (COVID-19) Outcomes: A Meta-Analysis. *Cureus* 2024;16:e59591.
- 82 Gaiha SM, Cheng J, Halpern-Felsher B. Association Between Youth Smoking, Electronic Cigarette Use, and COVID-19. *J Adolescent Health* 2020;67:519–23.
- 83 Molino AR, Jerry-Fluker J, Atkinson MA, *et al.* The association of alcohol, cigarette, e-cigarette, and marijuana use with disease severity in adolescents and young adults with pediatric chronic kidney disease. *Pediatr Nephrol* 2021;36:2493–7.
- 84 Ashour O, Al-Huneidy L, Noordeen H. The implications of vaping on surgical wound healing: A systematic review. *Surgery* 2023;173:1452–62.
- 85 Agochukwu N, Liao JY. Debunking the myth of e-cigarettes: A case of free flap compromise due to e-cigarette use within the first 24 hours. *J Plast Reconstr Aesthet Surg* 2018;71:451–3.
- 86 Kandel ER, Kandel DB. A Molecular Basis for Nicotine as a Gateway Drug. *N Engl J Med* 2014;371:932–43.
- 87 Selya AS, Foxon F. Trends in electronic cigarette use and conventional smoking: quantifying a possible “diversion” effect among US adolescents. *Addiction* 2021;116:1848–58.
- 88 Vanyukov MM, Tarter RE, Kirillova GP, *et al.* Common liability to addiction and “gateway hypothesis”: theoretical, empirical and evolutionary perspective. *Drug Alcohol Depend* 2012;123 Suppl 1:S3–17.
- 89 St Helen G, Eaton DL. Public Health Consequences of e-Cigarette Use. *JAMA Intern Med* 2018;178:984–6.
- 90 Kaleta D, Wojtyński P, Polańska K. Use of electronic cigarettes among secondary and high school students from a socially disadvantaged rural area in Poland. *BMC Public Health* 2016;15:703.
- 91 Fergusson DM, Boden JM, Horwood LJ. The developmental antecedents of illicit drug use: Evidence from a 25-year longitudinal study. *Drug Alcohol Depend* 2008;96:165–77.
- 92 Kopa-Stojak PN, Pawliczak R. Comparison of the effects of active and passive smoking of tobacco cigarettes, electronic nicotine delivery systems and tobacco heating products on the expression and secretion of oxidative stress and inflammatory response markers. A systematic review. *Inhal Toxicol* 2024;36:75–89.
- 93 Li X, Yuan L, Wang F, *et al.* Health outcomes of electronic cigarettes. *Chin Med J* 2024;137:1903–11.
- 94 Rebuli ME, Rose JJ, Noël A, *et al.* The E-cigarette or Vaping Product Use-Associated Lung Injury Epidemic: Pathogenesis, Management, and Future Directions: An Official American Thoracic Society Workshop Report. *Ann Am Thorac Soc* 2023;20:1–17.
- 95 Sund LJ, Dargan PI, Archer JRH, *et al.* E-cigarette or vaping-associated lung injury (EVALI): a review of international case reports from outside the United States of America. *Clin Toxicol (Phila)* 2023;61:91–7.
- 96 MHRA. E-cigarette use or vaping: reporting suspected adverse reactions, including lung injury. 2020. Available: <https://www.gov.uk/drug-safety-update/e-cigarette-use-or-vaping-reporting-suspected-adverse-reactions-including-lung-injury>
- 97 Jongenelis MI, Gill M, Lawrence N, *et al.* Quitting intentions and behaviours among young Australian e-cigarette users. *Addiction* 2024;119:1608–15.
- 98 Lin C, Gaiha SM, Halpern-Felsher B. Nicotine Dependence from Different E-Cigarette Devices and Combustible Cigarettes among US Adolescent and Young Adult Users. *IJERPH* 2022;19:5846.
- 99 Cuccia AF, Patel M, Amato MS, *et al.* Quitting e-cigarettes: Quit attempts and quit intentions among youth and young adults. *Prev Med Rep* 2021;21:101287.
- 100 World Health Organization. WHO report on the global tobacco epidemic 2019: offer help to quit tobacco use. 2019. Available: <https://www.who.int/publications/i/item/9789241516204>
- 101 Graham-DeMello A, Sloan O, Frost K, *et al.* Young people's experiences of addiction to nicotine vaping products: A qualitative analysis from Aotearoa New Zealand. *Drug Alcohol Rev* 2024;43:1710–7.
- 102 Al-Hamdani M, Hopkins DB, Hardardottir A, *et al.* Perceptions and Experiences of Vaping Among Youth and Young Adult E-Cigarette Users: Considering Age, Gender, and Tobacco Use. *J Adolesc Health* 2021;68:787–93.

- 103 Leslie FM. Unique, long-term effects of nicotine on adolescent brain. *Pharmacol Biochem Behav* 2020;197:173010.
- 104 Castro EM, Lotfipour S, Leslie FM. Nicotine on the developing brain. *Pharmacol Res* 2023;190:106716.
- 105 Elatfy A, Vrahimis S, Conti A, *et al.* Chronic tobacco smoking and neurocognitive impairments in adolescents and young adults: a systematic review and meta-analysis. *Front Psychiatry* 2024;15:1384408.
- 106 Gogliettino AR, Potenza MN, Yip SW. White matter development and tobacco smoking in young adults: A systematic review with recommendations for future research. *Drug Alcohol Depend* 2016;162:26–33.
- 107 Tobore TO. On the potential harmful effects of E-Cigarettes (EC) on the developing brain: The relationship between vaping-induced oxidative stress and adolescent/young adults social maladjustment. *J Adolesc* 2019;76:202–9.
- 108 Augenstein JA, Smaldone AM, Usseglio J, *et al.* Electronic Cigarette Use and Academic Performance Among Adolescents and Young Adults: A Scoping Review. *Acad Pediatr* 2024;24:228–42.
- 109 Glantz SA, Nguyen N, Oliveira da Silva AL. Population-Based Disease Odds for E-Cigarettes and Dual Use versus Cigarettes. *NEJM Evid* 2024;3:EVIDoa2300229.
- 110 Li L, Asemota I, Liu B, *et al.* AMSTAR 2 appraisal of systematic reviews and meta-analyses in the field of heart failure from high-impact journals. *Syst Rev* 2022;11:147.
- 111 Lorenz RC, Matthias K, Pieper D, *et al.* AMSTAR 2 overall confidence rating: lacking discriminating capacity or requirement of high methodological quality? *J Clin Epidemiol* 2020;119:142–4.
- 112 Bojicic R, Todoric M, Puljak L. Most systematic reviews reporting adherence to AMSTAR 2 had critically low methodological quality: a cross-sectional meta-research study. *J Clin Epidemiol* 2024;165:111210.
- 113 Georges Khalil DM, Ribeiro N, Jantz M, *et al.* A systematic review of symptom clusters and common diagnostic results related to lung injuries due to vaping among adolescents and young adults prospero 2020 crd42020185520. 2020. Available: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020185520
- 114 Carmen María Suárez Freire JRB. Prevalence of respiratory symptoms in young adults who use electronic cigarettes: a systematic review. prospero 2024 crd42024527302. 2024. Available: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42024527302
- 115 Andre Duraes MF, Neves J. Respiratory symptoms associated with the use of electronic nicotine delivery systems in adolescents and young adults: a systematic review. prospero 2024 crd42024553301. 2024. Available: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42024553301
- 116 Ali Ziyab DZ, Albatineh A. Association between electronic cigarette use and asthma symptoms among young people: a systematic review and meta-analysis. prospero 2020 crd42020180452. 2020. Available: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020180452
- 117 Li J, Hui X, Lu Z, *et al.* Association between initial use of e-cigarettes and subsequent cigarette smoking among adolescents and young adults: a systematic review and meta-analysis prospero 2020 crd420201632422. n.d. Available: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD420201632422
- 118 Ilaria Alabiso CB, Haw S, Whittaker AA. The relationship between e-cigarette uptake and subsequent substance use is a current matter of debate in public health. is e-cigarette 'ever use' associated with substance use and/or alcohol use in young people aged 11-19? prospero 2021 crd42021244724. 2021. Available: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42021244724
- 119 Stephanie Clendennen MH, Mason E, Khan J, *et al.* Does e-cigarette use predict marijuana use among youth and young adults? a systematic review and meta-analysis prospero 2021 crd42021276586. 2021. Available: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42021276586
- 120 Matheus Fernandez FGD, Cascaes AM, Muniz F. Electronic nicotine delivery systems and dental caries in adolescents and adults: a systematic review with meta-analysis. 2023. Available: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42023478845
- 121 Gaiha S, Halpern-Felsher B. Assessing the dual burden of adolescent e-cigarette use and depression and reduction strategies, 2020. Available: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020161803
- 122 Erick Santana de Souza LRdM, Aldair Jose de Oliveira. The effects of e-cigarette use on physical activity levels in adolescents: a systematic review and meta-analysis. 2023. Available: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42023442372
- 123 Luiz Felipe Frazão DR. The risks of using electronic cigarettes among young people: a systematic review. 2023. Available: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42023467968
- 124 Yee Yau Lee KZT, Wong MHH, Mahdi SS, *et al.* The use of e-cigarettes in young adults - a systematic review of the risk-benefit analysis. 2024. Available: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42024536596
- 125 Conde M, Tudor K, Begh R, *et al.* Electronic cigarettes and subsequent use of cigarettes in young people: An evidence and gap map. *Addiction* 2024;119:1698–708.